

ADAPTED COMPLEX TUBES ON THE SYMPLECTIZATION OF PSEUDO-HERMITIAN MANIFOLDS

GIUSEPPE TOMASSINI AND SERGIO VENTURINI

ABSTRACT. Let (M, θ) be a pseudo-Hermitian space of real dimension $2n + 1$, that is M is a CR-manifold of dimension $2n + 1$ and θ is a contact form on M giving the Levi distribution $HT(M) \subset TM$. Let $M^\theta \subset T^*M$ be the canonical symplectization of (M, θ) and M be identified with the zero section of M^θ . Then M^θ is a manifold of real dimension $2(n + 1)$ which admit a canonical foliation by surfaces parametrized by $\mathbb{C} \ni t + i\sigma \mapsto \phi_p(t + i\sigma) = \sigma \theta_{g_t(p)}$, where $p \in M$ is arbitrary and g_t is the flow generated by the Reeb vector field associated to the contact form θ .

Let J be an (integrable) complex structure defined in a neighbourhood U of M in M^θ . We say that the pair (U, J) is an adapted complex tube on M^θ if all the parametrizations $\phi_p(t + i\sigma)$ defined above are holomorphic on $\phi_p^{-1}(U)$.

In this paper we prove that if (U, J) is an adapted complex tube on M^θ , then the real function E on $M^\theta \subset T^*M$ defined by the condition $\alpha = E(\alpha)\theta_{\pi(\alpha)}$, for each $\alpha \in M^\theta$, is a canonical equation for M which satisfies the homogeneous Monge-Ampère equation $(dd^c E)^{n+1} = 0$.

We also prove that if M and θ are real analytic then the symplectization M^θ admits an unique maximal adapted complex tube.

1. ADAPTED COMPLEX TUBES

In this paper we apply some results obtained by the authors in [7] to the study of suitable “adapted” integrable complex structure on open neighbourhood of a pseudo-Hermitian manifold in its symplectization.

We follow [2] for standard notations in complex and CR-geometry.

Given a smooth real differentiable manifold M of dimension m we denote by T^*M its cotangent bundle and $\pi : T^*M \rightarrow M$ the canonical projection.

Let θ be a smooth 1-form on M such that $\theta_p \neq 0$ for each $p \in M$. Then the subset of by T^*M

$$M^\theta = \{ \alpha \in T^*M \mid \alpha \wedge \theta_{\pi(\alpha)} = 0 \},$$

is the line bundle on $M \subset T^*M$ of the forms α which are proportional to the form $\theta_{\pi(\alpha)}$. We denote by $E : M^\theta \rightarrow \mathbb{R}$ the function which satisfies

$$\alpha = E(\alpha)\theta_{\pi(\alpha)}$$

2000 *Mathematics Subject Classification.* Primary 32W20, 32V40 Secondary 53D10, 35Fxx.

Key words and phrases. Complex manifolds, CR Manifolds, Complex Monge-Ampère equation, Contact geometry.

for each $\alpha \in T^*M$.

Denoting by M_0 the zero section of T^*M we see that for each $\alpha \in M^\theta$ we have $E(\alpha) = 0$ if, and only if, $\alpha \in M_0$.

We freely identify the manifold M with the zero section M_0 in M^θ .

It easy to show that M^θ is a smooth closed submanifold of T^*M of (real) dimension $m + 1$ and E is a smooth function on M .

We also denote by X^θ the vector field on T^*M which is the infinitesimal generator of the one parameter group of transformation defined by

$$h_t^\theta(\alpha) = \alpha + t\theta_{\pi(\alpha)}$$

for each $\alpha \in T^*M$ and $t \in \mathbb{R}$.

Let $x = (x_1, \dots, x_m)$ be a local coordinate system on M and let $(x, p) = (x_1, \dots, x_m, p_1, \dots, p_m)$ the corresponding local coordinate system on T^*M , so that

$$\theta = \sum_{i=1}^m p_i(\theta) dx_i.$$

Then we have

$$X^\theta = \sum_{i=1}^m p_i(\theta) \frac{\partial}{\partial p_i}.$$

Clearly the vector field X^θ is tangent to M^θ and $X^\theta(E) = 1$. Thus the smooth function E has no critical points on M^θ .

Let now (M, θ) be a *pseudo-Hermitian manifold*, that is M is a orientable CR-manifold of dimension $2n + 1$ with non degenerate Levi form and θ is a non degenerate real 1-form which vanishes on the Levi distribution $HT(M) \subset TM$ and defines the *pseudo-Hermitian structure* on M . Then (M, θ) is a contact manifold with volume form $\theta \wedge (d\theta)^n$ and there exists a vector field ξ_0^θ , the *Reeb vector field*, which is characterized by the conditions $\xi_0^\theta \lrcorner \theta = 1$ and $\xi_0^\theta \lrcorner d\theta = 0$. The pull-back of $d\theta$ is a symplectic form on M^θ that will be called the *symplectization* of M .

Let g_t be the one parameter (local) group of transformations associated to ξ_0^θ , that is for each $p \in M$ the map

$$g_t(p) : I_p \rightarrow M, \quad t \mapsto g_t(p)M, \quad I_p \subset \mathbb{R}$$

is the maximal integral curve of ξ_0^θ such that $g_0(p) = p$.

For each $p \in M$ let $\tilde{I}_p = \{z = t + i\sigma \in \mathbb{C} \mid t \in I_p\}$ and the map

$$\phi_p = \phi_p^\theta : \tilde{I}_p \rightarrow M^\theta$$

defined by the formula

$$\phi_p^\theta(z) = \phi^\theta(t + i\sigma) = \sigma \theta_{g_t(p)}.$$

We also denote by ξ^θ the vector field on M^θ which is the infinitesimal generator of the one parameter local group of transformations defined by

$$g_t^\theta(\alpha) = E(\alpha) \theta_{g_t(p)},$$

where $p = \pi(\alpha)$.

Then the vector field ξ^θ extends ξ_0^θ and it is easy to show that the one parameter groups g_t^θ and h_t^θ commute and generate a two dimensional distribution on M^θ whose leaves are parametrized by the maps ϕ_p^θ . Moreover we have

$$(1) \quad \begin{aligned} \xi^\theta(E) &= 0, \\ X^\theta(E) &= 1, \\ [\xi^\theta, X^\theta] &= 0. \end{aligned}$$

We now give the main definition of this paper.

Definition 1.1. Let (M, θ) be a pseudo-Hermitian manifold of real dimension $2n + 1$. Let the manifold M be identified with the zero section M_0 in M^θ .

An adapted complex tube on M^θ is a pair (U, J) where U is a open subset of M^θ containing the zero section M_0 and J is an (integrable) complex structure on U which satisfies the following conditions:

- (1) for each $p \in M$ the set $U_p = \{\alpha \in U \mid \pi(\alpha) = p\}$ is connected;
- (2) the restriction of J to the Levi distribution $HT(M)$ is the given CR structure of $HT(M)$;
- (3) if $p \in M$ the restriction of the map ϕ_p^θ to the open set $\tilde{I}_p \cap (\phi_p^\theta)^{-1}(U)$ is holomorphic.

An adapted complex tube (U, J) with J of class C^k , $k = 1, \dots, \infty, \omega$, is said to be C^k -maximal, if for any other adapted complex tube on M^θ such that J' is of class C^k , $U \subset U'$ and $J|_U = J'$ one has $U = U'$ (and hence $J' = J$).

Adapted tubes of class C^∞ or C^ω are said to be smooth or real analytic, respectively.

The main results of this paper are the following.

Theorem 1.1. Let (M, θ) be a pseudo-Hermitian manifold of real dimension $2n + 1$ with M and θ of class C^∞ . Let (U, J) be any smooth adapted complex tube on M^θ . Then the function E satisfies

$$(2) \quad \left\{ \begin{aligned} &(\ddot{d}d^c E)^{n+1} = 0 \text{ on } U, \\ &dE \wedge d^c E \wedge (\ddot{d}d^c E)^n \neq 0 \text{ near } M, \\ &E|_M = 0, \\ &d^c E|_{T(M)} = -\theta. \end{aligned} \right.$$

Theorem 1.2. Let (M, θ) be a pseudo-Hermitian manifold of real dimension $2n + 1$ with M and θ real analytic. Then there exists a unique real analytic maximal adapted complex tube (U, J) on M^θ .

Remark 1.1. Adapted complex structure on the tangent (and cotangent) bundle a Riemannian manifolds are studied in [5], [6] and, independently, in [4].

2. PROOF OF THEOREM 1.1

Let (M, θ) be a fixed pseudo-Hermitian manifold.

Lemma 2.1. *If (U, J) is any smooth adapted complex tube on M^θ then*

$$(3) \quad J(\xi^\theta) = X^\theta.$$

Proof. Let $\alpha \in U \subset M^\theta$ and $p = \pi(\alpha)$. Then $\alpha = \sigma_0 \theta_p$, where $\sigma_0 = E(\alpha)$. By definition of ξ^θ

$$\begin{aligned} \xi^\theta(\alpha) &= \left. \frac{d}{dt} g_t^\theta(\alpha) \right|_{t=0} = \left. \frac{d}{dt} (\sigma_0 \theta_{g_t(p)}) \right|_{t=0} \\ &= \left. \frac{d}{dt} \phi_p(t + i\sigma_0) \right|_{t=0}. \end{aligned}$$

Since the map $\phi_p(t + i\sigma)$ is holomorphic we obtain

$$\begin{aligned} J(\xi^\theta(\alpha)) &= \left. \frac{d}{d\sigma} \phi_p(i(\sigma_0 + \sigma)) \right|_{\sigma=0} = \left. \frac{d}{d\sigma} (\alpha + \sigma \theta_p) \right|_{\sigma=0} \\ &= \left. \frac{d}{d\sigma} h_\sigma^\theta(\alpha) \right|_{\sigma=0} = X^\theta(\alpha), \end{aligned}$$

as desired.

□

Using the terminology of [7] we observe that by (1) and (3) the pair $(\xi^\theta, -E)$ is a smooth *calibrated foliation*, that is

$$\begin{aligned} [\xi^\theta, J(\xi^\theta)] &= 0, \\ dE(\xi^\theta) &= 0, \\ d^c E(\xi^\theta) &= -1. \end{aligned}$$

Moreover the function E vanishes on M and the vector field ξ^θ extends the Reeb vector field ξ_0^θ on M . Since the Reeb vector field ξ_0^θ is an infinitesimal symmetry for the contact distribution $HT(M)$ from of [7, Theorem 3.1] it follows that the form $\xi^\theta \lrcorner dd^c E = L_{\xi^\theta} d^c E$ vanishes identically on U . Here $L_{\xi^\theta} d^c E$ stands for the Lie derivative of the form $d^c E$ with respect to the vector field ξ^θ . Since $\xi^\theta \neq 0$ on U it follows that the form $dd^c E$ satisfies the Monge-Ampère equation $(dd^c E)^{n+1} = 0$ on U .

Clearly, the function E vanishes exactly on M and hence the restriction of $d^c E$ to TM is $\lambda \theta$ for some (smooth) function $\lambda : M \rightarrow \mathbb{R}$. For each $p \in M$ we have

$$\begin{aligned} \lambda(p) &= \lambda(p) \theta_p(\xi_0^\theta(p)) = d^c E(\xi_0^\theta(p)) \\ &= d^c E(\xi^\theta(p)) = -dE(X^\theta(p)) = -1 \end{aligned}$$

and hence $d^c E|_{T(M)} = -\theta$.

Since $\theta \wedge (d\theta)^n$ is a volume form on M we have $dE \wedge d^c E \wedge (dd^c E)^n \neq 0$ in a neighbourhood of M .

This completes the proof of Theorem 1.1 .

3. PROOF OF THEOREM 1.2

Let (M, θ) be a fixed pseudo-Hermitian manifold.

Let $\Phi : M \times \mathbb{R} \rightarrow M^\theta$ be the map defined by $\Phi(p, \sigma) = \sigma \theta_p$ for each $p \in M$ and each $\sigma \in \mathbb{R}$. Then Φ is a diffeomorphism between $M \times \mathbb{R}$ and M^θ .

By [1] there exists a pair $(\tilde{M}, j)$ where $\tilde{M}$ is a complex manifold and j is a real analytic embedding of M in $\tilde{M}$. Then, by analytic extension, there exist an open neighbourhood V of $M \times \{0\}$ in $M \times \mathbb{C}$ and a map $\Psi^c : V \rightarrow \tilde{M}$ satisfying

- (1) for each $p \in M$ the set $I_p = \{t \in \mathbb{R} \mid (p, t) \in V\}$ is an open interval in $\mathbb{R}$ and $\Psi^c(p, t) = g_t(p)$ is the maximal integral curve of the vector field ξ_0^θ such that $g_0(p) = 0$, i.e. $\Psi^c(p, 0) = 0$ for each $p \in M$;
- (2) for each $(p, t) \in V \cap M \times \mathbb{R}$ the set $V_{(p, t)} = \{\sigma \in \mathbb{R} \mid (p, t + i\sigma) \in V\}$ is an open interval in $\mathbb{R}$ containing the origin $0 \in \mathbb{R}$
- (3) for each $p \in M$ the map $z = (t + i\sigma) \mapsto \Psi^c(p, z)$ is holomorphic when defined.

Then, set $W = \{(p, \sigma) \in M \times \mathbb{R} \mid (p, i\sigma) \in V\}$ and define $\Psi : W \rightarrow \tilde{M}$ by the formula $\Psi(p, \sigma) = \Psi^c(p, i\sigma)$. Since $X^\theta(p) = J(\xi_0^\theta(p))$ is transversal to M for each $p \in M$ it follows that, shrinking V (and hence W) and $\tilde{M}$ if necessary, the map Ψ is a real analytic diffeomorphism between W and $\tilde{M}$.

Let $U = \Phi(V)$ and J be the pullback under $\Phi \circ \Psi^{-1}$ of the complex structure of $\tilde{M}$. It is then easy to show that (U, J) is a real analytic adapted complex tube on M^θ .

This proves the existence of real analytic adapted complex tubes.

Let now (U_1, J_1) and (U_2, J_2) be two real analytic adapted complex tubes on M^θ . We claim that J_1 and J_2 agree on the intersection $U_1 \cap U_2$. By an analytic continuation argument it suffices to prove that J_1 and J_2 agree on a neighbourhood of M in $U_1 \cap U_2$.

Again by [1], there exist open neighbourhoods $V_1 \subset U_1$ and $V_2 \subset U_2$ of M and a real analytic diffeomorphism $F : V_1 \rightarrow V_2$ which is the identity map on M and which is holomorphic when V_1 and V_2 are endowed by the complex structures J_1 and J_2 respectively .

We are going to prove that $V_1 = V_2$ and the map F actually is the identity map. Indeed, let $\alpha \in V_1$ be arbitrary, $p = \pi(\alpha)$ and $\sigma_0 = E(\alpha)$. Then $\alpha = \sigma_0 \theta_p$. The maps $f_1(z) = \phi_p(z)$ and $f_2(z) = F(\phi_p(z))$ are both holomorphic and coincide when z is real. By analytic continuation $f_1 = f_2$. In particular, we have

$$F(\alpha) = F(\phi_p(i\sigma_0)) = \phi_p(i\sigma_0) = \alpha,$$

that is the map F is the identity map on V_1 , as claimed.

Now a standard argument implies that there exist a unique real analytic complex structure J on the set U defined as the union of all the open neighbourhood U' of M which admit a complex structure J' which make the pair (U', J') an adapted complex tube on M^θ .

The pair (U, J) is then the required unique real analytic maximal adapted complex tube for M^θ .

This concludes the proof of Theorem 1.2.

Remark 3.1. The construction of the map Ψ is similar to a one given in [3].

REFERENCES

- [1] A. Andreotti and G. A. Fredricks. Embeddability of Real Analytic Cauchy-Riemann Manifolds. *Annali della Scuola Normale Superiore*, 6:297–301, 1979.
- [2] S. Dragomir and G. Tomassini. *Differentiable Geometry and Analysis on CR Manifolds*. Birkhauser, 2006.
- [3] T. Duchamp and M. Kalka. Singular Monge-Ampère foliations. *Math. Ann.*, 325:187–209, 2003.
- [4] V. Guillemin and M. Stenzel. Grauert tubes and the homogeneous Monge-Ampère equation. *Journal of Differential Geometry*, 34:561–570, 1991.
- [5] L. Lempert and R. Szöke. Global solutions of the homogeneous complex Monge-Ampère equation and complex structures on the tangent bundle of Riemannian manifolds. *Math. Ann.*, 290:689–712, 1991.
- [6] R. Szöke. Complex structures on tangent bundles of Riemannian manifolds. *Math. Ann.*, 291:409–428, 1991.
- [7] G. Tomassini and S. Venturini. Contact geometry of one dimensional holomorphic foliations. *arXiv:0907.5082v1*, pages 1–15, 2009.

G. TOMASSINI: SCUOLA NORMALE SUPERIORE, PIAZZA DEI CAVALIERI, 7 — I-56126 PISA, ITALY

E-mail address: g.tomassini@sns.it

S. VENTURINI: DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA S. DONATO 5 — I-40127 BOLOGNA, ITALY

E-mail address: venturin@dm.unibo.it